

Comment on the new $D_s^{(*)+}\pi^0$ resonances

Thomas E. Browder,¹ Sandip Pakvasa,¹ and Alexey A. Petrov^{2,3}

¹*Department of Physics and Astronomy, University of Hawaii at Manoa, Honolulu, Hawaii 96822*

²*Department of Physics and Astronomy, Wayne State University, Detroit, MI 48201*

³*Michigan Center for Theoretical Physics, University of Michigan, Ann Arbor, MI 48109*

We propose an explanation of the new resonances observed in $D_s^{(*)+}\pi^0$ decays. We suggest that the data can be explained by the mixing of conventional p-wave excited D_s^+ mesons with 4-quark states. The narrow states observed in $D_s^+\pi^0$ and $D_s^{*+}\pi^0$ are primarily p-wave D_{sJ}^* states, while the predominantly 4-quark states are shifted above $D^{(*)}K$ threshold and should be broad. Ranges for the mixing parameter and mass of the 4-quark state in this scenario are given. Other experimental consequences are discussed.

PACS numbers: 12.20.He, 13.25.Hw, 13.60.Rj

Meson spectroscopy is an important laboratory for understanding quark confinement. Mesons containing one heavy quark can provide invaluable information about the structure of the QCD Lagrangian, as spectroscopic considerations simplify significantly in the limit of infinitely heavy quark. In this limit the heavy quark spin S_Q decouples, so the total angular momentum of the light degrees of freedom J_l^P becomes a “good” quantum number. This leads to an important prediction of heavy quark symmetry: the appearance of heavy meson states in the form of degenerate parity doublets classified by the total angular momentum of the light degrees of freedom. This mass degeneracy is lifted with the inclusion of subleading $1/m_Q$ corrections. This useful picture is built into many quark-model descriptions of heavy meson spectra. The resulting models have been very successful in explaining the spectrum of negative-parity scalar and vector $J_l^P = 1/2^-$ and positive-parity vector and tensor $J_l^P = 3/2^+$ states.

A narrow resonance in $D_s^+\pi^0$ was recently reported by BaBar [1] and confirmed by the CLEO [2] and Belle [3] collaborations. Its decay patterns suggest a quark-model 0^+ classification, which would identify it with the positive-parity $J_l^P = 1/2^+$ p-wave state. As in the D meson system, p-wave states for the D_s^+ system are expected, and two narrow states, $D_{s1}(2536)$ and $D_{s2}(2573)$ were discovered by ARGUS and CLEO, respectively [4]. In analogy to the D system, two broad states are also expected.

The mass of the new state 2317.6 ± 1.3 MeV appears surprisingly low for quark model practitioners. In fact, this state appears below DK threshold, closing off the most natural decay channel for this state. This forces it to decay mainly via an isospin-violating transition into the $D_s^+\pi^0$ final state and makes its width quite narrow. Its mass disagrees with most existing predictions of quark models [5, 6, 7, 8, 9, 10]. For example, a mass of 2487 MeV is obtained in the recent potential model calculation by Eichten and Di Pierro [6]. Quenched lattice calculations also seem to favor larger values of the mass of this state [11] (see, however, [12]). This led to a lively discussion of the possible non- $q\bar{q}$ nature of this state [13, 14, 15, 16]. In addition, a second narrow state is observed in $D_s^{*+}\pi^0$ at a mass near 2460 MeV [2, 3]. This state would naturally be identified as a spin 1 positive parity p-wave meson. However, its mass also appears too low for the potential model expectations (e.g. 2605 MeV [6]).

The observed low values of the masses for these states, however, do not signal a breakdown of quark-model descriptions of the heavy meson spectrum, as it is difficult to assess the accuracy of these predictions, especially in the charm sector. Many authors make use of the non-relativistic nature of the charm quark, taking into account $1/m_c$ corrections only. For the 0^+ state, quark model predictions range from the values of 2387 – 2395 MeV [7, 9] (still above the DK threshold) on the low end of the spectrum to 2508 MeV [10] on the high end. Since the described phenomena are highly non-perturbative, one should be careful before making a judgment on the nature of a given state based solely on the prediction of a given quark model. For example, as discussed above, in the $m_c \rightarrow \infty$ limit the 0^+ and 1^+ states are expected to become degenerate in mass, $m_{0^+}, m_{1^+} \rightarrow M$. This can be emulated in quark models by neglecting heavy-quark symmetry-violating $1/m_c$ corrections. Yet, different quark models predict very different behavior in this “heavy-quark limit”: for instance, one potential model [7] predicts that the mass M of the $(0^+, 1^+)$ multiplet will decrease to approximately 2382 MeV (which is less than the mass of the 0^+ state predicted in this model with the full

potential), while in a QCD string model [9] it is expected to increase up to 2500 MeV (which is much greater than the mass of the 0^+ state predicted in this model with the full potential). In addition, quark models, modified to include chiral symmetry constraints, generally predicted lower values of mass splitting between $(0^-, 1^-)$ and $(0^+, 1^+)$ multiplets, of the order of 200 – 300 MeV [17]. Nevertheless, the fact that the new state appears below DK threshold and is almost degenerate with a non-strange 0^+ p -wave D state [18] is curious and deserves an investigation.

It has been shown that the proper description of the low lying scalar mesons requires the inclusion of the coupling of the scalar $q\bar{q}$ states to the s -wave meson-meson channel, and this coupling is very important [19]. By an extension of this argument, we would expect the scalar 0^+ $c\bar{s}$ state to mix with the DK s -wave state near threshold. This mixing can occur quite naturally if the states have the same quantum numbers (for instance, via the common intermediate states, see Fig. 1) and could shift the mass of the $c\bar{s}$ state.

Alternatively, the possibility of 4-quark states with quark content $c\bar{s}q\bar{q}$ has been discussed in the past [20, 21]. Here we consider the possibility that there is a 4-quark state that mixes with the $c\bar{s}$ state and shifts its mass below DK threshold.




FIG. 1: Example of the interaction generating mixing of the $q\bar{q}$ and 4-quark states. In general, all possible (multibody) intermediate states such as $D^{(*)}K^{(*)}$, $DK\pi\pi$, ... will contribute.

We briefly review two-state mixing for the case of two nearby hadronic states. Let m_0 be the unshifted mass of the $q\bar{q}$ state and $\tilde{m}_0$ be the unshifted mass of the 4-quark state. The strong interaction can mix these states by introducing non-diagonal matrix elements δ in the mass matrix of $qq\bar{q}\bar{q}$ and $q\bar{q}$ states, which has to be diagonalized. The masses of the two mixed (mass eigen)states, m_1 and m_2 will then be given by

$$m_1 = \frac{1}{2}\Sigma + \frac{1}{2}\sqrt{\Delta^2 + 4\delta^2} \quad (1)$$

$$m_2 = \frac{1}{2}\Sigma - \frac{1}{2}\sqrt{\Delta^2 + 4\delta^2} \quad (2)$$

where $\Sigma = m_0 + \tilde{m}_0$ and $\Delta = m_0 - \tilde{m}_0$. The mixing angle that defines the composition of resulting mass eigenstates in terms of the original $qq\bar{q}\bar{q}$ and $q\bar{q}$ states is given by $\tan(2\theta) = 2\delta/\Delta$.

Let us be more specific. For example, take $\tilde{m}_0$ the bare mass of the 4-quark state to be just above DK threshold at 2.37 GeV and m_0 at the value 2.48 GeV as given by the potential model; then for a mixing parameter δ of 0.092 GeV, we find two mixed states with masses $m_2 = 2.3194$ GeV and $m_1 = 2.5375$ GeV. The corresponding mixing angle is 28.8° . The lower state, which is dominantly a p -wave D_s meson, is below DK threshold, 2.358 GeV, but above $D\pi$ threshold, 2.103 GeV. Conversely, the higher state, which is dominantly a 4-quark system, is above DK threshold and is broad. There is a large set of parameters for the bare 4 quark mass and mixing that can shift the mass of the D_{sJ}^* to the observed value of 2317 GeV. This is illustrated in Figure 2. If $\tilde{m}_0$ is too high, it is difficult to shift the D_{sJ}^* state, with a reasonable amount of mixing.

In principle, we also expect the same type of mixing to occur for the 1^+ state. For instance, a molecular-type bound-state of D^*K of the kind suggested by Lipkin [13, 20] and others, can mix with the $c\bar{s}$ 3P_1 state and shift the 1^+ states as well. To calculate the allowed range of mixing in this case is more complicated since there is mixing between the two 1^+ states in addition to the mixing between the lower 1^+ state and the relevant 4 quark state, and we do not attempt a detailed fit.

The properties and decay modes of the lower mixed states should be very similar to those expected for p -wave D_s mesons. Like the broad D^{**} states perhaps some of the additional states may be identified in exclusive B decays such as $B \rightarrow D^{(*)0}\bar{D}^0K$ with enough statistics. The observation of these extra states could lend extra support for the existence of the multi-quark and/or molecular states.

The 4 quark states can also give rise to very distinctive final states that are doubly charged [20, 21]. For example, there may be $I = 1$ states including a doubly charged state and a neutral state with decay modes into D^+K^+ , etc. In the scenario described here, these exotic final states are above DK threshold and are therefore quite broad, experimentally difficult to find or even may be non-existent as "real" bound states. Note that we do not expect narrow doubly charged $D_s^+\pi^+$ or $D_s^+\pi^-$ states in the scenarios described above.

It has also been suggested [13, 22] that radiative transitions of the type $D_{sJ}^*(2317) \rightarrow D_s^*\gamma$ could be used to test the nature of these states. In particular, it was suggested that a very small branching ratio for the radiative decays would be an indication of a non- $q\bar{q}$ nature of these states. In the scheme described above this test will be useful only for very small or very large (about $\pi/2$) values of the mixing angle between $c\bar{s}$ and $c\bar{s}q\bar{q}$ states, as the contribution of $c\bar{s}q\bar{q} \rightarrow D_s^*\gamma$ transition is suppressed by $\sin\theta$. This is not so for moderate values of the mixing angle [23].




FIG. 2: δ versus m_0 for solutions that satisfy the constraints of this scenario.

In conclusion, we propose that the new states observed in $D_s^+\pi^0$ and $D_s^{*+}\pi^0$ are shifted by mixing with 4-quark states. We expect that the observed states will have properties similar to those expected for p -wave D_s mesons except for their masses. The additional states responsible for the mixing and mass shifts will be above $D^{(*)}K$ threshold and have large widths.

We thank M. Suzuki for useful discussions. We acknowledge support from the U.S. Department of Energy under grants DE-FG03-94ER40833, DE-FG02-96ER41005.

-
- [1] B. Aubert *et al.* (BaBar Collaboration), hep-ex/030402, Phys. Rev. Lett. **90**, 242001 (2003).
 - [2] D. Besson *et al.* (CLEO Collaboration), hep-ex/0305100, Phys. Rev. **D 68**, 032002 (2003).
 - [3] K. Abe *et al.* (Belle Collaboration), (hep-ex/0307052), submitted to Phys. Rev. Lett. ; K. Abe *et al.* (Belle Collaboration), (hep-ex/0308019) to appear in Phys. Rev. Lett.

- [4] D.E. Groom *et al.* (Particle Data Group), Eur. Phys. J. C **15**, 1 (2000).
- [5] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985).
- [6] M. Di Pierro and E. Eichten, Phys. Rev. D **64**, 114004 (2001).
- [7] S. N. Gupta and J. M. Johnson, Phys. Rev. D **51**, 168 (1995).
- [8] J. Zeng, J. W. Van Orden and W. Roberts, Phys. Rev. D **52**, 5229 (1995).
- [9] Y. S. Kalashnikova, A. V. Nefediev and Y. A. Simonov, Phys. Rev. D **64**, 014037 (2001).
- [10] D. Ebert, V. O. Galkin and R. N. Faustov, Phys. Rev. D **57**, 5663 (1998) [Erratum-ibid. D **59**, 019902 (1999)].
- [11] R. Lewis and R. M. Woloshyn, Phys. Rev. D **62**, 114507 (2000); J. Hein *et al.*, Phys. Rev. D **62**, 074503 (2000); G. S. Bali, hep-ph/0305209.
- [12] A. Dougall, R. D. Kenway, C. M. Maynard and C. McNeile, arXiv:hep-lat/0307001.
- [13] T. Barnes, F. E. Close and H. J. Lipkin, arXiv:hep-ph/0305025.
- [14] H. Y. Cheng and W.-S. Hou, arXiv:hep-ph/0305038.
- [15] A. P. Szczepaniak, arXiv:hep-ph/0305060.
- [16] R.N. Cahn and J.D. Jackson, arXiv:hep-ph/0305012; Y. B. Dai, C. S. Huang, C. Liu and S. L. Zhu, arXiv:hep-ph/0306274.
- [17] W. A. Bardeen and C. T. Hill, Phys. Rev. D **49**, 409 (1994); W. A. Bardeen, E. J. Eichten and C. T. Hill, arXiv:hep-ph/0305049; D. Ebert, T. Feldmann and H. Reinhardt, Phys. Lett. B **388**, 154 (1996); D. Ebert, T. Feldmann, R. Friedrich and H. Reinhardt, Nucl. Phys. B **434**, 619 (1995); M. A. Nowak, M. Rho and I. Zahed, Phys. Rev. D **48**, 4370 (1993); M. A. Nowak, M. Rho and I. Zahed, arXiv:hep-ph/0307102; A. Deandrea *et al.*, Phys. Rev. D **58**, 034004, (1998) [arXiv:hep-ph/9802308]; A. Deandrea, G. Nardulli, A.D. Polosa, arXiv:hep-ph/0307069.
- [18] K. Abe *et al.* (Belle Collaboration), arXiv:hep-ex/0307021, submitted to Phys. Rev. D.
- [19] E. van Beveren *et al.*, Z. Phys. C **30**, 615 (1986); E. van Beveren and G. Rupp, hep-ph/0304105; E. van Beveren and G. Rupp, Phys. Rev. Lett. **91**, 012003 (2003); J. Weinstein and N. Isgur, Phys. Rev. D **41**, 2236 (1990).
- [20] H.J. Lipkin, Phys. Lett. B. **70**, 113 (1977).
- [21] M. Suzuki and S.F. Tuan, Phys. Lett. B. **133**, 125 (1983).
- [22] S. Godfrey, arXiv:hep-ph/0305122; P. Colangelo and F. De Fazio, arXiv:hep-ph/0305140.
- [23] T. E. Browder, S. Pakvasa and A. A. Petrov, in preparation.